



# POST REHABILITATION IMPACT ON SYRIAN REFUGEES WITH LOWER LIMB AMPUTATION

ALIREZA HEIDARI  
RELIEF INTERNATIONAL, USA

19<sup>th</sup> International Conference on  
**OTOLARYNGOLOGY: ENT SURGERY**

October 04-05, 2022 | Berlin, Germany

**T**here is evidence on the deterioration of smell and taste in patients with COVID-19. Differences were found in the prevalence of these symptoms in different populations. Based on the pattern of presentation, as of April 16, 2020, the definition of a suspected case in Argentina incorporated these symptoms. The objective of this study is to establish the prevalence of these deficits in patients with COVID-19.

**Methods:** Multicenter study. Confirmed COVID-19 cases older than 18 years, RT-PCR swabs, between May 15 and August 31, 2020 participated. Participants completed an online survey, which included sociodemographic questions, on COVID-19 symptoms, comorbidities and in relation to perception of loss of smell and taste.

**Results:** 241 patients participated, 79% reported loss of smell and 54% of taste. In 32% of the cases the loss of smell was the first symptom. The proportion of men and women who reported olfactory dysfunction was similar and the same happened in relation to the loss of taste. The prevalence of these symptoms coincides in relation to the age of the patient.

**Discussion:** A high proportion of patients with COVID-19 present olfactory and taste dysfunction in the Argentine population. The dissemination of the olfactory and gustatory test adapted to cultural smells, which can be carried out at home, is important. In the future we will develop the implementation of a massive disposable and digitized test.

**Methods:** Multicenter study. Confirmed COVID-19 cases older than 18 years, RT-PCR swabs, between May 15 and August 31, 2020 participated. Participants completed an online survey, which included sociodemographic questions, on COVID-19 symptoms, comorbidities and in relation to perception of loss of smell and taste.

**Results:** 241 patients participated, 79% reported loss of smell and 54% of taste. In 32% of the cases the loss of smell was the first symptom. The proportion of men and women who reported olfactory dysfunction was similar and the same happened in relation to the loss of taste. The prevalence of these symptoms coincides in relation

**Discussion:** A high proportion of patients with COVID-19 present olfactory and taste dysfunction in the Argentine population. The dissemination of the olfactory and gustatory test adapted to cultural smells, which can be carried out at home, is important. In the future we will develop the implementation of a massive disposable and digitized test

COFRE A has completed his Medicine Degree in 2004 at the School of Medicine of the University of Buenos Aires, Argentina. She is a teacher at the Medical Career at the Alarez Hospital in Buenos Aires. He is currently serving as president of the Asociación Red Olfato Argentina & Smell Training, carrying out a vigorous research task with another look at otorhinolar here is evidence on the deterioration of smell and taste in patients with COVID-19. Differences were found in the prevalence of these

symptoms in different populations. Based on the pattern of presentation, as of April 16, 2020, the definition of a suspected case in Argentina incorporated these symptoms. The objective of this study is to establish the prevalence of these deficits in patients with COVID-19.

Methods: Multicenter study. Confirmed COVID-19 cases older than 18 years, RT-PCR swabs, between May 15 and August 31, 2020 participated. Participants completed an online survey, which included sociodemographic questions, on COVID-19 symptoms, comorbidities and in relation to perception of loss of smell and taste.

Results: 241 patients participated, 79% reported loss of smell and 54% of taste. In 32% of the cases the loss of smell was the first symptom. The proportion of men and women who reported olfactory dysfunction was similar and the same happened in relation to the loss of taste. The prevalence of these symptoms coincides in relation to the age of the patient.

Discussion: A high proportion of patients with COVID-19 present olfactory and taste dysfunction in the Argentine population. The dissemination of the olfactory and gustatory test adapted to cultural smells, which can be carried out at home, is important. In the future we will develop the implementation of a massive disposable and digitized test.

## Speaker Biography

COFRE A has completed his Medicine Degree in 2004 at the School of Medicine of the University of Buenos Aires, Argentina. She is a teacher at the Medical Career at the Alarez Hospital in Buenos Aires. He is currently serving as president of the Asociación Red Olfato Argentina & Smell Training, carrying out a vigorous research task with another look at otorhinolaryngology. Member of different scientific societies. here is evidence on the deterioration of smell and taste in patients with COVID-19. Differences were found in the prevalence of these symptoms in different populations. Based on the pattern of presentation, as of April 16, 2020, the definition of a suspected case in Argentina incorporated these symptoms. The objective of this study is to establish the prevalence of these deficits in patients with COVID-19.

COFRE A has completed his Medicine Degree in 2004 at the School of Medicine of the University of Buenos Aires, Argentina. She is a teacher at the Medical Career at the Alarez Hospital in Buenos Aires. He is currently serving as president of the Asociación Red Olfato Argentina & Smell Training, carrying out a vigorous research task with another look at otorhinolaryngology. Member of different scientific societies. here is evidence on the deterioration of smell and taste in patients with COVID-19. Differences were found in the prevalence of these symptoms in different populations. Based on the pattern of presentation, as of April 16, 2020, the definition of a suspected case in Argentina incorporated these symptoms. The objective of this study is to establish the prevalence of these deficits in patients with COVID-19.

## References

1. Mercer PE, Woodcock HV, Eley JD, Platé M, Sulikowski MG, et al. (2016) Exploration of a potent PI3 kinase/mTOR inhibitor as a novel anti-fibrotic agent in IPF. *Thorax* 71: 701-711.
2. Behr J, Neuser P, Prasse A, Kreuter M, Rabe K, et al. (2017) Exploring efficacy and safety of oral Pirfenidone for progressive, non-IPF lung fibrosis (RELIEF)-a randomized, double-blind, placebo-controlled, parallel group, multi-center, phase II trial. *BMC pulmonary medicine* 17 : 122.
3. Dhooria S, Agarwal R, Gupta D (2015) Is pirfenidone ready for use in non-idiopathic pulmonary fibrosis interstitial lung diseases? *Lung India: official organ of Indian Chest Society* 32:4.
4. Bendstrup E, Maher TM, Manali ED, Wijsenbeek M (2015) Challenges in the classification of fibrotic ILD. *Sarcoidosis vasculitis and diffuse lung disease* 32: 4-9.
5. Yamakawa H, Kitamura H, Takemura T, Ikeda S, Sekine A (2018) Prognostic factors and disease behaviour of pathologically proven fibrotic non-specific interstitial pneumonia. *Respirology* 23: 1032-1040
6. Brown JC, Kraft J, Vega-Olivo M, Galli J, Simpson S. et al. (2017) Rate of decline in fvc in non-ipf pulmonary fibrosis after anti-fibrotic initiation. *Inc38. American Thoracic Society A5402-A5402*.
7. Nair GB, Matela A, Kurbanov D, Raghu G (2016) Newer developments in idiopathic pulmonary fibrosis in the era of anti-fibrotic medications. *Expert review of respiratory medicine* 10: 699-711.
8. Wells AU, Brown KK, Flaherty KR, Kolb M, Thannickal VJ (2018) what's in a name? That which we call IPF, by any other name would act the same. *ER Journal* 51: 1800692
9. Tzouveleakis A, Bouros D (2020) Endotyping of progressive fibrotic interstitial lung diseases: It is the final destination that matters and not the journey. *EBioMedicine* 1:51
10. Ryerson CJ, Urbania TH, Richeldi L, Mooney JJ, Lee JS, et al. (2013) Prevalence and prognosis of unclassifiable interstitial lung disease. *ERJ*

11. Jo HE, Corte TJ, Moodley Y, Levin K, Westall G. Evaluating the interstitial lung disease multidisciplinary meeting: A survey of expert centres. *BMC pulmonary medicine* 16: 22
12. Adegunsoye A, Streck ME. Therapeutic approach to adult fibrotic lung diseases (2016) *Chest* 150: 1371-1386.
13. Datta A, Scotton CJ, Chambers RC (2011) Novel therapeutic approaches for pulmonary fibrosis. *British j pharmacology* 163: 141-712.
14. Prognostic factors and disease behaviour of pathologically proven fibrotic non-specific interstitial pneumonia. *Respirology* 23: 1032-1040
15. Brown JC, Kraft J, Vega-Olivo M, Galli J, Simpson S. et al. (2017) Rate of decline in fvc in non-ipf pulmonary fibrosis after anti-fibrotic initiation. *Inc38. American Thoracic Society A5402-A5402*.
16. Nair GB, Matela A, Kurbanov D, Raghu G (2016) Newer developments in idiopathic pulmonary fibrosis in the era of anti-fibrotic medications. *Expert review of respiratory medicine* 10: 699-711.
17. Wells AU, Brown KK, Flaherty KR, Kolb M, Thannickal VJ (2018) what's in a name? That which we call IPF, by any other name would act the same. *ER Journal* 51: 1800692
18. Tzouvelekis A, Bouros D (2020) Endotyping of progressive fibrotic interstitial lung diseases: It is the final destination that matters and not the journey. *EBioMedicine* 1:51
19. Ryerson CJ, Urbania TH, Richeldi L, Mooney JJ, Lee JS, et al. (2013) Prevalence and prognosis of unclassifiable interstitial lung disease. *ERJ* 42: 750-757